

Badly approximable numbers related to the Littlewood conjecture

Moshchevitin N.G.¹

Abstract.

By means of Peres-Schlag's method we prove the existence of real numbers α, β such that

$$\liminf_{q \rightarrow \infty} (q \log^2 q) \|\alpha q\| \|\beta q\| > 0.$$

1. The famous Littlewood conjecture suppose that for any two real numbers α, β one has

$$\liminf_{q \rightarrow \infty} q \|\alpha q\| \|\beta q\| = 0$$

(here $\|\cdot\|$ denotes the distance to the nearest integer).

This conjecture is obviously true for *almost all* (in the sense of the Lebesgue measure) pairs $(\alpha, \beta) \in \mathbb{R}^2$. Moreover, from Gallagher's theorem (see [1]) we know that for almost all α, β one has

$$\liminf_{q \rightarrow \infty} (q \log^2 q) \|\alpha q\| \|\beta q\| = 0.$$

Here we prove the following result.

Theorem 1. *There exist real numbers α, β such that*

$$\liminf_{q \rightarrow \infty} (q \log^2 q) \|\alpha q\| \|\beta q\| > 0.$$

Our result is based on an original method of constructing badly approximable numbers invented by Y. Peres and W. Schlag in [2]. This method was recently applied by the author to several Diophantine problems [3]-[6]. In the sequel we suppose that we deal with the dyadic logarithm $\log x = \log_2 x$.

2. In this section we prove an elementary lemma dealing with the sum

$$\sigma(\alpha; q, Q) = \sum_{x=q}^{Q-1} \frac{1}{\|\alpha x\| x \log_2^2 x}.$$

We take $\alpha \in \mathbb{R}$ to be a badly approximable number. This means that with some positive $\delta < 1$ one has

$$\inf_{q \in \mathbb{N}} q \|\alpha q\| \geq \delta. \quad (1)$$

Lemma 1. *Let (1) holds. Then for any natural numbers $Q > q$ one has*

$$\sigma(\alpha; q, Q) \leq 2^8 \delta^{-1} \left(\log_2 \left(\frac{\log_2 Q}{\log_2 q} \right) + \log_2(1/\delta) \right).$$

¹ Research is supported by grants RFFI 06-01-00518

Proof. For natural μ, ν define

$$A_{\mu, \nu} = \{x \in \mathbb{N} : 2^\nu \leq x < 2^{\nu+1}, 2^{-\mu-1} < \|\alpha x\| \leq 2^{-\mu}\}.$$

As for $2^\nu \leq x < 2^{\nu+1}$ from (1) follows

$$\delta 2^{-\nu-1} < \|\alpha x\| \leq 2^{-1},$$

we see that

$$\{x \in \mathbb{N} : 2^\nu \leq x < 2^{\nu+1}\} = \bigsqcup_{\mu=1}^{\nu+1+\lceil \log_2(1/\delta) \rceil} A_{\mu, \nu}.$$

Now

$$\begin{aligned} \sigma(\alpha; q, Q) &\leq \sum_{\nu=\lfloor \log_2 q \rfloor}^{\lfloor \log_2 Q \rfloor} \sum_{x=2^\nu}^{2^{\nu+1}-1} \frac{1}{\|\alpha x\| x \log_2^2 x} = \sum_{\nu=\lfloor \log_2 q \rfloor}^{\lfloor \log_2 Q \rfloor} \sum_{\mu=1}^{\nu+1+\lceil \log_2(1/\delta) \rceil} \sum_{x \in A_{\mu, \nu}} \frac{1}{\|\alpha x\| x \log_2^2 x} \leq \\ &\leq \sum_{\nu=\lfloor \log_2 q \rfloor}^{\lfloor \log_2 Q \rfloor} \sum_{\mu=1}^{\nu+1+\lceil \log_2(1/\delta) \rceil} 2^{\mu-\nu+1} \nu^{-2} \text{card}(A_{\mu, \nu}). \end{aligned}$$

Note that

$$\text{card}(A_{\mu, \nu}) \leq \text{card}(\{x \in \mathbb{N} : x < 2^{\nu+1}, \|\alpha x\| \leq 2^{-\mu}\}) = I_{\mu, \nu}.$$

We have

$$\sigma(\alpha; q, Q) \leq \sum_{\nu=\lfloor \log_2 q \rfloor}^{\lfloor \log_2 Q \rfloor} \sum_{\mu=1}^{\nu+1+\lceil \log_2(1/\delta) \rceil} 2^{\mu-\nu+1} \nu^{-2} I_{\mu, \nu}. \quad (2)$$

The value of $I_{\mu, \nu}$ is bounded by the number of integer points in the region

$$\Omega = \{(x, y) \in \mathbb{R}^2 : 0 \leq x < 2^{\nu+1}, |x\alpha - y| \leq 2^{-\mu}\}.$$

Obviously the Lebesgue measure of Ω is equal to $\text{mes}\Omega = 2^{\nu-\mu+1}$.

Let there exists an integer primitive point $(p_0, a_0) \in \Omega$ (otherwise $I_{\mu, \nu} = 0$). From (1) we deduce

$$\delta p_0^{-1} \leq \|p_0 \alpha\| \leq 2^{-\mu}$$

and $p_0 \geq \delta 2^\mu$. Now we consider two cases.

In the *first case* we suppose that (p_0, a_0) is the unique primitive integer point in $\Omega(p)$. Then

$$I_{\mu, \nu} \leq \lfloor 2^{\nu+1}/p_0 \rfloor + 1 \leq 2^{\nu-\mu+2} \delta^{-1}.$$

In the *second case* the convex hull of all integer points in the region Ω is a convex polygon Π with integer vertices. According to Pick's formula the number $I_{\mu, \nu}$ of integer point in the polygon Π is not greater than $6\text{mes}\Pi$ and so

$$I_{\mu, \nu} \leq 6\text{mes}\Pi \leq 6\text{mes}\Omega = 12 \cdot 2^{\nu-\mu}.$$

In any case

$$I_{\mu, \nu} \leq 2^{\nu-\mu+4} \delta^{-1}. \quad (3)$$

Now we substitute (3) into (2) and obtain

$$\begin{aligned}\sigma(\alpha; q, Q) &\leq 2^{5\delta-1} \cdot \sum_{\nu=\lfloor \log_2 q \rfloor}^{\lfloor \log_2 Q \rfloor} (\nu + 1 + \lceil \log_2(1/\delta) \rceil) \nu^{-2} \leq 2^{6\delta-1} \left(\sum_{\nu=\lfloor \log_2 q \rfloor}^{\lfloor \log_2 Q \rfloor} \nu^{-1} + \log_2(1/\delta) \right) \leq \\ &\leq 2^{8\delta-1} \left(\log_2 \left(\frac{\log_2 Q}{\log_2 q} \right) + \log_2(1/\delta) \right)\end{aligned}$$

and Lemma 1 is proved.

3. In this section we (following Y. Peres and W. Schlag [2]) construct "dangerous" sets of reals. Let ε be positive and small enough. For integers $2 \leq x, 0 \leq y \leq x$ define

$$E(x, y) = \left[\frac{y}{x} - \frac{\varepsilon}{\|\alpha x\| x^2 \log_2^2 x}, \frac{y}{x} + \frac{\varepsilon}{\|\alpha x\| x^2 \log_2^2 x} \right], \quad E(x) = \bigcup_{y=0}^x E(x, y) \cap [0, 1]. \quad (4)$$

Define

$$l_0 = 0, \quad l_x = \lfloor \log_2(\|\alpha x\| x^2 \log_2^2 x / 2\varepsilon) \rfloor, \quad x \in \mathbb{N}. \quad (5)$$

Each segment from the union $E_\alpha(x)$ from (4) can be covered by a dyadic interval of the form

$$\left(\frac{b}{2^{l_x}}, \frac{b+z}{2^{l_x}} \right), \quad z = 1, 2.$$

Let $A(x)$ be the smallest union of all such dyadic segments which cover the whole set $E(x)$. Put

$$A^c(x) = [0, 1] \setminus A(x).$$

Then

$$A^c(x) = \bigcup_{\nu=1}^{\tau_x} I_\nu$$

where closed segments I_ν are of the form

$$\left[\frac{a}{2^{l_x}}, \frac{a+1}{2^{l_x}} \right], \quad a \in \mathbb{Z}. \quad (6)$$

We take q_0 to be a large positive integer. In order to prove Theorem 1 it is sufficient to show that for all $q \geq q_0$ the sets

$$B_q = \bigcap_{x=q_0}^q A^c(x)$$

are not empty. Indeed as the sets B_q are closed and nested we see that there exists real β such that

$$\beta \in \bigcap_{q \geq q_0} B_q.$$

One can see that the pair α, β satisfies the conclusion of Theorem 1.

In the next section we prove the following statement.

Lemma 2. *Suppose that ε is small enough. Then for q_0 large enough and for any*

$$q_1 \geq q_0, \quad q_2 = q_1^3, \quad q_3 = q_2^3$$

the following holds. If

$$\text{mes} B_{q_2} \geq \text{mes} B_{q_1}/2 > 0 \quad (7)$$

then

$$\text{mes} B_{q_3} \geq \text{mes} B_{q_2}/2 > 0. \quad (8)$$

Theorem 1 follows from Lemma 2 by induction as the base of the induction obviously follows from the arguments of Lemma's proof.

4. In this section we prove Lemma 2. Here we also follow the arguments from the paper [2] by Y. Peres and W. Schlag. First of all we show that for $x \geq q^3$ where $q \geq q_0$ one has

$$\text{mes} \left(B_q \cap A(x) \right) \leq \frac{2^4 \varepsilon}{\|\alpha x\| x \log_2^2 x} \times \text{mes} B_q. \quad (9)$$

Indeed as from (5) it follows that

$$l_x \leq l = \lfloor \log_2(q^2 \log_2^2 q / \varepsilon) \rfloor, \quad \forall x \leq q$$

we see that B_q is a union

$$B_q = \bigcup_{\nu=1}^{T_q} J_\nu$$

with J_ν of the form

$$\left[\frac{a}{2^l}, \frac{a+1}{2^l} \right], \quad a \in \mathbb{Z}.$$

Note that $A(x)$ consists of the segments of the form (6) and for $x \geq q^3 > 2^{l+1}$ (for q_0 large enough) we see that each J_ν has at least two rational fractions of the form $\frac{y}{x}, \frac{y+1}{x}$ inside. So

$$\text{mes}(J_\nu \cap A(x)) \leq \frac{2^4 \varepsilon}{\|\alpha x\| x \log_2^2 x} \times \text{mes} J_\nu. \quad (10)$$

Now (9) follows from (10) by summation over $1 \leq \nu \leq T_q$.

To continue we observe that

$$B_{q_3} = B_{q_2} \setminus \left(\bigcup_{x=q_2+1}^{q_3} A(x) \right),$$

and hence

$$\text{mes} B_{q_3} \geq \text{mes} B_{q_2} - \sum_{x=q_2+1}^{q_3} \text{mes}(B_{q_2} \cap A(x)).$$

As

$$B_{q_2} \cap A(x) \subseteq B_{q_1} \cap A_{\alpha_i}(x)$$

we can apply (9) for every x from the interval $q_1^3 \leq q_2 < x \leq q_3$:

$$\text{mes}(B_{q_2} \cap A(x)) \leq \text{mes}(B_{q_1} \cap A(x)) \leq \frac{2^4 \varepsilon}{\|\alpha x\| x \log_2^2 x} \times \text{mes} B_{q_1} \leq \frac{2^5 \varepsilon}{\|\alpha x\| x \log_2^2 x} \times \text{mes} B_{q_2}$$

(in the last inequality we use the condition (7) of Lemma 2). Now as $\frac{\log_2 q_3}{\log_2 q_2} = 3$ the conclusion (8) of Lemma 2 follows from Lemma 1 for ε small enough:

$$\begin{aligned} \text{mes} B_{q_3} &\geq \text{mes} B_{q_2} (1 - 2^5 \varepsilon \sigma(\alpha; q_2 + 1, q_3 + 1)) \geq \\ &\geq \text{mes} B_{q_2} \left(1 - 2^{14} \varepsilon \left(\log_2 \left(\frac{\log_2 q_3}{\log_2 q_2} \right) + \log_2(1/\delta) \right) \right) \geq \text{mes} B_{q_2}/2. \end{aligned}$$

References

- [1] Gallagher P. Metric simultaneous Diophantine approximations. // J. London Math. Soc. 37 (1962), p. 387 – 390.
- [2] Peres Y., Schlag W. Two Erdős problems on lacunary sequences: chromatic numbers and Diophantine approximations. // Preprint, available at: arXiv: 0706.0223v1 [math.CO] 1Jun2007.
- [3] Moshchevitin N.G. A version of the proof for Peres-Schlag's theorem on lacunary sequences. // Preprint, available at arXiv: 0708.2087v2 [math.NT] 15Aug2007
- [4] Moshchevitin N.G. Density modulo 1 of sublacunary sequences: application of Peres-Schlag's arguments. // Preprint, available at arXiv: 0709.3419v2 [math.NT] 20Oct2007
- [5] Moshchevitin N.G. On small fractional parts of polynomials. // J. Number Theory (2008), doi:10.1016/j.jnt.2008.07.008
- [6] Moshchevitin N.G. Towards BAD conjecture. // Preprint, available at arXiv: 0712.2423v2 12Apr2008.

author: Nikolay Moshchevitin

e-mail: moshchevitin@mech.math.msu.su, moshchevitin@rambler.ru